

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 108 435 A1

(12)

EUROPEAN PATENT APPLICATION
published in accordance with Art. 158(3) EPC

(43) Date of publication:
20.06.2001 Bulletin 2001/25

(51) Int Cl.7: **A61K 45/00**, A61K 39/395

(21) Application number: 99938591.7

(86) International application number:
PCT/JP99/04533

(22) Date of filing: 23.08.1999

(87) International publication number:
WO 00/10607 (02.03.2000 Gazette 2000/09)

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE**
Designated Extension States:
AL LT LV MK RO SI

(72) Inventors:
• **FUNAKOSHI, Akihiro**
Fukuoka-shi, Fukuoka 815-0042 (JP)
• **MIYASAKA, Kyoko**
Setagaya-ku, Tokyo 156-0042 (JP)

(30) Priority: 24.08.1998 JP 25179698
02.03.1999 JP 5430299

(74) Representative: Geering, Keith Edwin
REDDIE & GROSE
16 Theobalds Road
London WC1X 8PL (GB)

(71) Applicant: **CHUGAI SEIYAKU KABUSHIKI
KAISHA**
Tokyo, 115-8543 (JP)

(54) **PREVENTIVES OR REMEDIES FOR PANCREATITIS CONTAINING IL-6 ANTAGONISTS AS
THE ACTIVE INGREDIENT**

(57) A preventive or therapeutic agent for pancrea-
titis said agent comprising as an active ingredient an in-

terleukin-6 (IL-6) antagonist such as an antibody direct-
ed against IL-6 receptor.

EP 1 108 435 A1

Description

Technical Field

[0001] The present invention relates to a preventive or therapeutic agent for pancreatitis comprising an interleukin-6 (IL-6) antagonist as an active ingredient.

Background Art

[0002] IL-6 is a cytokine which is also called B cell stimulating factor 2 (BSF2) or interleukin β 2. IL-6 was discovered as a differentiation factor involved in the activation of B-lymphatic cells (Hirano, T. et al., Nature (1986) 324, 73-76). Thereafter, it was found to be a multifunctional cytokine that influences various functions of cells (Akira, S. et al., Adv. in Immunology (1993) 54, 1-78). IL-6 has been reported to induce the maturation of T-lymphatic cells (Lotz, M. et al., J. Exp. Immunol. (1988) 167, 1253-1258).

[0003] IL-6 transmits its biological activity through two types of proteins on the cell. One is IL-6 receptor, a ligand-binding protein with a molecular weight of about 80 kD, to which IL-6 binds (Taga, T. et al., J. Exp. Med. (1987) 166, 967-981; Yamasaki, K. et al., Science (1987) 241, 825-828). IL-6 receptor occurs not only in the membrane-bound form that penetrates through and is expressed on the cell membrane but also as a soluble IL-6 receptor consisting mainly of the extracellular region.

[0004] The other is a membrane-bound protein gp130 having a molecular weight of about 130 kD that is involved in non-ligand-binding signal transduction. IL-6 and IL-6 receptor form the IL-6/IL-6 receptor complex, which, after binding to gp130, transmits its biological activity to the cell (Taga, T. et al., Cell (1989) 58, 573-581).

[0005] An IL-6 antagonist is a substance that inhibits the transduction of biological activity of IL-6. As the IL-6 antagonist, there have been known so far antibody directed against IL-6 (anti-IL-6 antibody), antibody directed against IL-6 receptor (anti-IL-6 receptor antibody), and antibody directed against gp130 (anti-gp130 antibody), altered IL-6, partial peptides of IL-6 or IL-6 receptor and the like.

[0006] Anti-IL-6 receptor antibody has been described in several reports (Novick D. et al., Hybridoma (1991) 10, 137-146, Huang, Y. W. et al., Hybridoma (1993) 12, 621-630, International Patent Publication WO 95-09873, French Patent Application FR 2694767, United States Patent US 521628). Humanized PM-1 antibody has been known that was obtained by grafting the complementarity determining region (CDR) of one of them, a mouse antibody PM-1 (Hirata, Y. et al., J. Immunology (1989) 143, 2900-2906), to a human antibody (the International Patent Publication WO 92-19759).

[0007] Pancreatitis is an inflammatory disease in which the activation of pancreatic enzymes causes autolysis in pancreatic tissues. There have been reported

that the amount of IL-6 produced in the peripheral blood mononuclear cells is significantly high in patients with pancreatitis as compared to healthy normal humans, and that IL-6 production from the peripheral blood mononuclear cells is high in cases of acute pancreatitis with systemic complications as compared to those with no complications (de Beaux A. C. et al., Brit. J. Surgery, 83, 1071-5, 1996). Furthermore, since blood levels of IL-6 are higher and respond earlier than other parameters in severe cases of acute pancreatitis, they have been considered to be a prognostic indicator for severity of pancreatitis (Inagaki, T. et al., Pancreas, 14, 1-8, 1997).

[0008] It has been suggested that IL-1 and TNF closely correlate with the disease states, and mice lacking receptors to both of the cytokines do not suffer serious disease conditions, and show markedly decreased mortality rate (Denham, W. et al., Gastroenterology, 113, 1741-6, 1997). Attempts have been made to treat pancreatitis using these inhibitors in animal models (Norman, J. et al., Surgery, 117, 648-6755, 1995, Hughes, C. B. et al., American J. Surgery, 171, 274-280, 1996, Norman, J. et al., surgery, 120, 515-621, 1996).

[0009] However, no attempts have been made to specifically suppress the biological activity of IL-6 using IL-6 antagonists such as anti-IL-6 receptor antibody in pancreatitis, and it was unknown that IL-6 antagonists such as anti-IL-6 receptor antibody exhibit therapeutic effects on pancreatitis.

Disclosure of the Invention

[0010] It is an object of the present invention to provide a preventive or therapeutic agent for pancreatitis said agent being free of the above-mentioned drawbacks.

[0011] Thus, the present invention provides (1) a preventive or therapeutic agent for pancreatitis comprising an IL-6 antagonist as an active ingredient.

[0012] The present invention also provides (2) a preventive or therapeutic agent for pancreatitis comprising an antibody directed against IL-6 receptor as an active ingredient.

[0013] The present invention also provides (3) a preventive or therapeutic agent for pancreatitis comprising a monoclonal antibody directed against IL-6 receptor as an active ingredient.

[0014] The present invention also provides (4) a preventive or therapeutic agent for pancreatitis comprising a monoclonal antibody directed against human IL-6 receptor as an active ingredient. The monoclonal antibody directed against human IL-6 receptor is preferably PM-1 antibody.

[0015] The present invention also provides (5) a preventive or therapeutic agent for pancreatitis comprising a monoclonal antibody directed against mouse IL-6 receptor as an active ingredient. The monoclonal antibody directed against mouse IL-6 receptor is preferably

MR16-1 antibody.

[0016] The present invention also provides (6) a preventive or therapeutic agent for pancreatitis comprising a recombinant antibody directed against IL-6 receptor as an active ingredient. The recombinant antibody directed against IL-6 receptor has preferably a human antibody constant region (C region).

[0017] The present invention also provides (7) a preventive or therapeutic agent for pancreatitis comprising a chimeric or humanized antibody directed against IL-6 receptor as an active ingredient.

[0018] The present invention also provides (8) a preventive or therapeutic agent for pancreatitis comprising humanized PM-1 antibody as an active ingredient.

[0019] The present invention also provides (9) a preventive or therapeutic agent for acute or chronic pancreatitis comprising the IL-6 antagonist described in the above (1) to (8) as an active ingredient. Acute or chronic pancreatitis is, for example, severe or mild pancreatitis.

[0020] The present invention also provides (10) an agent for suppressing pancreatic edema, said agent comprising the IL-6 antagonist described in the above (1) to (8) as an active ingredient.

[0021] The present invention also provides (11) an agent for suppressing pancreatic edema, said agent comprising an antibody directed against IL-6 receptor described in the above (3) to (8) as an active ingredient.

Brief Description of the Drawings

[0022] Figure 1 shows that caerulein administration causes pancreatic edema resulting in the increased weight of the pancreas in IL-6 transgenic mice as compared to normal mice. It also shows that the above effect is suppressed by the administration of anti-IL-6 antibody.

[0023] Figure 2 is a micrograph of the pancreatic tissue of a normal mouse that developed acute pancreatitis by caerulein administration.

[0024] Figure 3 is a micrograph of the pancreatic tissue of an IL-6 transgenic mouse that developed acute pancreatitis by caerulein administration.

[0025] Figure 4 is a micrograph of the pancreatic tissue of an IL-6 transgenic mouse that developed acute pancreatitis by caerulein administration and that received MR16-1 as well. As compared to the caerulein-induced pancreatitis in the normal mouse in Figure 2, the caerulein-induced pancreatitis is aggravated (thus, enhanced edema in the interstitial tissue and enhanced infiltration in inflammatory cells) in the IL-6 transgenic mouse in Figure 3. In Figure 4, in contrast, the aggravation has been suppressed by the administration of an anti-IL-6 receptor antibody MR16-1.

[0026] Figure 5 shows that the increase in the weight of the pancreas in an IL-6 transgenic mouse induced by the administration of LPS and caerulein can be suppressed by the administration of an anti-IL-6 receptor antibody.

Best Mode for Carrying Out the Invention

[0027] IL-6 antagonists for use in the present invention may be of any origin, any kind, and any form, as long as they have a preventive or therapeutic effect for pancreatitis, or an effect of controlling pancreatic edema.

[0028] IL-6 antagonists block signal transduction by IL-6 and inhibit the biological activity of IL-6. Preferably, IL-6 antagonists have an activity of inhibiting the binding of any of IL-6, IL-6 receptor, and gp130. As the IL-6 antagonists, there can be mentioned preferably anti-IL-6 antibody, anti-IL-6 receptor antibody, anti-gp130 antibody, altered IL-6, altered soluble IL-6 receptor, a partial peptide of IL-6 or IL-6 receptor, and low molecular weight substances having the same activity as these.

[0029] Anti-IL-6 antibodies for use in the present invention can be obtained as polyclonal or monoclonal antibodies using a known method. As the anti-IL-6 antibodies for use in the present invention, monoclonal antibodies of, in particular, a mammalian origin, are preferred. Monoclonal antibodies of a mammalian origin include those produced by a hybridoma and those produced by a host which has been transformed with an expression vector containing genetically engineered antibody genes. These antibodies, via binding to IL-6, block the binding of IL-6 to IL-6 receptor and, thereby, block signal transduction of the biological activity of IL-6 into the cell.

[0030] Examples of such antibodies include MH166 (Matsuda et al., Eur. J. Immunol. (1988) 18, 951-956) and SK2 antibody (Sato, K. et al., The 21st Nihon Menekigakkai Soukai (General Meeting of the Japan Immunology Society), Academic Record (1991) 21, 166) and the like.

[0031] An anti-IL-6 antibody-producing hybridoma can be basically constructed using a known procedure as described below. Thus, IL-6 may be used as a sensitizing antigen and is immunized in the conventional method of immunization. The immune cells thus obtained are fused with known parent cells in the conventional cell fusion process, and then monoclonal antibody-producing cells are screened by the conventional screening method to prepare the desired hybridoma.

[0032] Specifically, anti-IL-6 antibody may be obtained in the following manner. For example, a human IL-6 for use as the sensitizing antigen to obtain antibody can be obtained using the IL-6 gene/amino acid sequence disclosed in Eur. J. Biochem (1987) 168, 543-550, J. Immunol. (1988) 140, 1534-1541, or Agr. Biol. Chem. (1990) 54, 2685-2688.

[0033] After a suitable host cell is transformed by inserting the IL-6 gene sequence into a known expression vector system, the IL-6 protein of interest is purified from the host cell or the culture supernatant thereof by a known method, and the purified IL-6 protein can be used as the sensitizing antigen. Alternatively, a fusion protein of the IL-6 protein and another protein may be used as the sensitizing antigen.

[0034] Anti-IL-6 receptor antibodies for use in the present invention can be obtained as polyclonal or monoclonal antibodies using a known method. As the anti-IL-6 antibodies for use in the present invention, monoclonal antibodies of, in particular, a mammalian origin, are preferred. Monoclonal antibodies of a mammalian origin include those produced by a hybridoma and those produced by a host which has been transformed with an expression vector containing genetically engineered antibody genes. The antibodies, via binding to IL-6 receptor, inhibit the binding of IL-6 to IL-6 receptor, and thereby block the transduction of the biological activity of IL-6 into the cell.

[0035] Examples of such antibodies include MR16-1 antibody (Tamura, T., et al., Proc. Natl. Acad. Sci. USA (1993) 90, 11924-11928), PM-1 antibody (Hirata, et al., J. Immunology (1989) 143, 2900-2906), or AUK12-20 antibody, AUK64-7 antibody or AUK146-15 antibody (International Patent Publication WO 92-19759), and the like. Among them, PM-1 antibody is most preferred.

[0036] Incidentally, the hybridoma cell line which produces PM-1 antibody has been internationally deposited under the provisions of the Budapest Treaty as PM-1 on July 10, 1990 with the National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology, of 1-3, Higashi 1-chome, Tsukuba city, Ibaraki pref., Japan, as FERM BP-2998. The hybridoma cell line which produces MR16-1 antibody has been internationally deposited under the provisions of the Budapest Treaty as MR16-1 on March 13, 1997 with the National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology, of 1-3, Higashi 1-chome, Tsukuba city, Ibaraki pref., Japan, as FERM BP-5875.

[0037] Hybridomas producing anti-IL-6 receptor monoclonal antibody can be basically prepared using a known procedure as described below. Thus, IL-6 receptor is used as a sensitizing antigen and is immunized according to the conventional method of immunization. The immune cells thus obtained are fused with known parent cells in the conventional cell fusion process, and then monoclonal antibody-producing cells may be screened by the conventional screening method to prepare the desired hybridoma.

[0038] Specifically, anti-IL-6 receptor antibody may be prepared in the following manner. For example, human IL-6 receptor used as the sensitizing antigen for obtaining antibody can be obtained using the IL-6 receptor gene sequence/amino acid sequence disclosed in European Patent Application EP 325474, and mouse IL-6 receptor can be obtained using that disclosed in Japanese Unexamined Patent Publication (Kokai) 3 (1991)-155795.

[0039] There are two types of IL-6 receptor proteins: IL-6 receptor expressed on the cell membrane, and IL-6 receptor detached from the cell membrane (soluble IL-6 receptor) (Yasukawa et al., J. Biochem. (1990) 108, 673-676). Soluble IL-6 receptor antibody is composed

substantially of the extracellular region of the IL-6 receptor bound to the cell membrane, and thereby is different from the membrane-bound IL-6 receptor in that the latter lacks the transmembrane region or both of the transmembrane region and the intracellular region. As the IL-6 receptor protein, any IL-6 receptor can be used, as long as it can be used a sensitizing antigen for production of the IL-6 receptor antibody for use in the present invention.

[0040] After the gene sequence of IL-6 receptor is inserted into a known expression vector system to transform an appropriate host cell, the desired IL-6 receptor protein may be purified from the host cell or a culture supernatant thereof using a known method. The purified IL-6 receptor protein thus purified may be used as the sensitizing antigen. Alternatively, cells that are expressing IL-6 receptor or a fusion protein of the IL-6 receptor protein and another protein may be used as the sensitizing antigen.

[0041] *E. coli* that has a plasmid pIBBSF2R containing cDNA encoding human IL-6 receptor has been internationally deposited under the provisions of the Budapest Treaty as HB101-pIBBSF2R on January 9, 1989 with the National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology, of 1-3, Higashi 1-chome, Tsukuba city, Ibaraki pref., Japan, as FERM BP-2232.

[0042] Anti-gp130 antibodies for use in the present invention can be obtained as polyclonal or monoclonal antibodies using a known method. As the anti-gp130 antibodies for use in the present invention, monoclonal antibodies of, in particular, a mammalian origin, are preferred. Monoclonal antibodies of a mammalian origin include those produced by a hybridoma and those produced by a host which has been transformed with an expression vector containing genetically engineered antibody genes. The antibodies, via binding to gp130, inhibit the binding of IL-6/IL-6 receptor complex to gp130, and thereby block the transduction of the biological activity of IL-6 into the cell.

[0043] Examples of such antibodies include AM64 antibody (Japanese Unexamined Patent Publication (Kokai) 3(1991)-219894), 4B11 antibody and 2H4 antibody (US 5571513), B-S12 antibody and B-P8 antibody (Japanese Unexamined Patent Publication (Kokai) 8 (1996)-291199).

[0044] A monoclonal antibody-producing hybridoma can be basically created using a known procedure as described below. Thus, gp130 may be used as a sensitizing antigen and is immunized in the conventional method of immunization. The immune cells thus obtained are fused with known parent cells in the conventional cell fusion process, and then the monoclonal antibody-producing hybridomas are screened by the conventional screening method to prepare the desired hybridoma.

[0045] Specifically, monoclonal antibody may be obtained in the following manner. For example, gp130

used as the sensitizing antigen for antibody generation can be obtained using the gp130 gene sequence/amino acid sequence disclosed in European Patent Application EP 411946.

[0046] After a suitable host cell is transformed by inserting the gp130 gene sequence into a known expression vector system, the gp130 protein of interest is purified from the host cell or from the culture supernatant thereof. The purified gp130 receptor protein can be used as the sensitizing antigen. Alternatively, a fusion protein of the gp130 protein and another protein may be used as the sensitizing antigen.

[0047] Though mammals to be immunized with the sensitizing antigen are not specifically limited, they are preferably selected in consideration of their compatibility with the parent cell for use in cell fusion. They generally include rodents such as mice, rats, hamsters and the like.

[0048] Immunization of animals with a sensitizing antigen is carried out using a known method. A general method, for example, involves the intraperitoneal or subcutaneous administration of a sensitizing antigen to the mammal. Specifically, a sensitizing antigen which has been diluted and suspended in an appropriate amount of phosphate buffered saline (PBS) or physiological saline etc. is mixed, as desired, with an appropriate amount of a common adjuvant, for example Freund's complete adjuvant. After being emulsified, it is preferably administered to a mammal several times every 4 to 21 days. Alternatively a suitable carrier may be used at the time of immunization of the sensitizing antigen.

[0049] After immunization and the confirmation of the increase in the desired antibody levels in the serum, the immune cells are taken out from the mammal and are subjected to cell fusion. Preferred immune cells subjected to cell fusion include in particular the spleen cells.

[0050] The mammalian myeloma cells as the other parent cells which are subjected to cell fusion with the above-mentioned immune cells preferably include various known cell lines such as P3X63Ag8.653 (Kearney, J. F. et al., J. Immunol. (1979) 123: 1548-1550), P3X63Ag8U.1 (Current Topics in Microbiology and Immunology (1978) 81: 1-7), NS-1 (Kohler, G. and Milstein, C., Eur. J. Immunol. (1976) 6: 511-519), MPC-11 (Margulies, D.H. et al., Cell (1976) 8: 405-415), SP2/0 (Shulman, M. et al., Nature (1978) 276: 269-270), FO (de St. Groth, S. F. et al., J. Immunol. Methods (1980) 35: 1-21), S194 (Trowbridge, I.S., J. Exp. Med. (1978) 148: 313-323), R210 (Galfre, G. et al., Nature (1979) 277: 131-133) and the like.

[0051] Cell fusion between the above immune cells and the myeloma cells may be essentially conducted in accordance with a known method such as that described in Milstein et al. (Kohler, G. and Milstein, C., Methods Enzymol. (1981) 73: 3-46).

[0052] More specifically, the above cell fusion is carried out in the conventional nutrient broth in the pres-

ence of, for example, a cell fusion accelerator. As the cell fusion accelerator, for example, polyethylene glycol (PEG), Sendai virus (HVJ) and the like may be used, and, in addition, an adjuvant such as dimethyl sulfoxide etc. may be added as desired to enhance the efficiency of fusion.

[0053] The preferred ratio of the immune cells and the myeloma cells to be used is, for example, 1 to 10 times more immune cells than the myeloma cells. Examples of culture media to be used for the above cell fusion include RPMI1640 medium and MEM culture medium suitable for the growth of the above myeloma cell lines, and the conventional culture medium used for this type of cell culture, and besides a serum supplement such as fetal calf serum (FCS) may be added.

[0054] In cell fusion, predetermined amounts of the above immune cells and the myeloma cells are mixed well in the above culture liquid, to which a PEG solution previously heated to about 37 °C, for example a PEG solution with a mean molecular weight of about 1000 to 6000, is added at a concentration of 30 to 60% (w/v) and mixed to obtain the desired fusion cells (hybridomas). Then, by repeating the sequential addition of a suitable culture liquid and centrifugation to remove the supernatant, cell fusion agents etc. which are undesirable for the growth of the hybridoma can be removed.

[0055] Said hybridoma is selected by culturing in the conventional selection medium, for example, the HAT culture medium (a culture liquid containing hypoxanthine, aminopterin, and thymidine). Culturing in said HAT culture medium is continued generally for a period of time sufficient to effect killing of the cells other than the desired hybridoma (non-fusion cells), generally several days to several weeks. The conventional limiting dilution method is conducted in which the hybridomas that produce the desired antibody are screened and monoclally cloned.

[0056] In addition to obtaining the above hybridoma by immunizing an animal other than the human with an antigen, it is also possible to sensitize human lymphocytes in vitro with desired antigen or desired antigen-expressing cells, and the resulting sensitized B lymphocytes are fused with a human myeloma cell for example U266, to obtain the desired human antibody having the activity of binding to desired antigen or desired antigen-expressing cells (see Japanese Post-examined Patent Publication (Kokoku) No. 1(1989)-59878). Furthermore, a transgenic animal having a repertoire of all human antibody genes is immunized with the antigen or the antigen-expressing cells to obtain the desired human antibody in the method described above (see International Patent Publication WO 93/12227, WO 92/03918, WO 94/02602, WO 94/25585, WO 96/34096 and WO 96/33735).

[0057] The monoclonal antibody-producing hybridomas thus constructed can be subcultured in the conventional culture liquid, or can be stored for a prolonged period of time in liquid nitrogen.

[0058] In order to obtain monoclonal antibodies from said hybridoma, a method can be used in which said hybridoma is cultured in the conventional method and the antibodies are obtained as the supernatant, or a method in which the hybridoma is administered to and grown in a mammal compatible with said hybridoma and the antibodies are obtained as the ascites. The former method is suitable for obtaining high-purity antibodies, whereas the latter is suitable for a large scale production of antibodies.

[0059] Specifically a hybridoma producing anti-IL-6 receptor antibody can be constructed using the method disclosed in Japanese Unexamined Patent Publication (Kokai) 3(1989)-139293. It can be conducted by a method in which the PM-1 antibody-producing hybridoma that was internationally deposited under the provisions of the Budapest Treaty as FERM BP-2998 on July 10, 1990 with the National Institute of Bioscience and Human Technology, Agency of Industrial Science and Technology, of 1-3, Higashi 1-chome, Tsukuba city, Ibaraki pref., Japan, is intraperitoneally injected to BALB/c mice (manufactured by CLEA Japan) to obtain the ascites from which the PM-1 antibody is purified, or a method in which said hybridoma is cultured in a suitable culture medium such as the RPMI1640 medium containing 10% bovine fetal serum and 5% MB-Condimed H1 (manufactured by Boehringer Mannheim), the hybridoma SFM medium (manufactured by GIBCO-BRL), the PFHM-II medium (manufactured by GIBCO-BRL) and the like, and the PM-1 antibody can be purified from the culture supernatant.

[0060] A recombinant antibody which was produced by the recombinant gene technology in which an antibody gene was cloned from the hybridoma and integrated into a suitable vector which was then introduced into a host can be used in the present invention as monoclonal antibody (see, for example, Borrebaeck C.A.K., and Larrick J.W. THERAPEUTIC MONOCLONAL ANTIBODIES, published in the United Kingdom by MACMILLAN PUBLISHERS LTD. 1990).

[0061] Specifically, mRNA encoding the variable region (V) of the desired antibody is isolated from antibody-producing cells such as a hybridoma. The isolation of mRNA is conducted by preparing total RNA using, for example, a known method such as the guanidine ultracentrifuge method (Chirgwin, J.M. et al., Biochemistry (1979) 18, 5294-5299), the AGPC method (Chomczynski, P. et al., Anal. Biochem. (1987) 162, 156-159), and then mRNA is purified from the total RNA using the mRNA Purification kit (manufactured by Pharmacia) and the like. Alternatively, mRNA can be directly prepared using the Quick Prep mRNA Purification Kit (manufactured by Pharmacia).

[0062] cDNA of the V region of antibody may be synthesized from the mRNA thus obtained using a reverse transcriptase. cDNA may be synthesized using the AMV Reverse Transcriptase First-strand cDNA Synthesis Kit and the like. Alternatively, for the synthesis and amplifi-

cation of cDNA, the 5'-Ampli FINDER RACE Kit (manufactured by Clontech) and the 5'-RACE method (Frohman, M.A. et al., Proc. Natl. Acad. Sci. USA (1988) 85, 8998-9002; Belyavsky, A. et al., Nucleic Acids Res. (1989) 17, 2919-2932) that employs polymerase chain reaction (PCR) may be used. The desired DNA fragment is purified from the PCR product obtained and may be ligated to vector DNA. Moreover, a recombinant vector is constructed therefrom and then is introduced into *E. coli* etc., from which colonies are selected to prepare the desired recombinant vector. The base sequence of the desired DNA may be confirmed by a known method such as the dideoxy method.

[0063] Once the DNA encoding the V region of the desired antibody has been obtained, it may be ligated to DNA encoding the constant region (C region) of the desired antibody, which is then integrated into an expression vector. Alternatively, the DNA encoding the V region of the antibody may be integrated into an expression vector which already contains DNA encoding the C region of the antibody.

[0064] In order to produce the antibody for use in the present invention, the antibody gene is integrated as described below into an expression vector so as to be expressed under the control of an expression regulatory region, for example an enhancer and/or a promoter. Subsequently, the expression vector may be transformed into a host cell and the antibody can then be expressed therein.

[0065] In accordance with the present invention, artificially altered recombinant antibody such as chimeric antibody and humanized antibody can be used for the purpose of lowering heterologous antigenicity against humans. These altered antibodies can be produced using known methods.

[0066] Chimeric antibody can be obtained by ligating the thus obtained DNA encoding the V region of antibody to DNA encoding the C region of human antibody, which is then integrated into an expression vector and introduced into a host for production of the antibody therein (see European Patent Application EP 125023, and International Patent Publication WO 92-19759). Using this known method, chimeric antibody useful for the present invention can be obtained.

[0067] For example, the plasmid that contains DNA encoding the L chain V region or the H chain V region of chimeric PM-1 antibody was designated as pPM-k3 or pPM-h1, respectively, and *E. coli* having the plasmid has been internationally deposited under the provisions of the Budapest Treaty as NCIMB 40366 and NCIMB 40362, respectively, on February 11, 1991 with the National Collections of Industrial and Marine Bacteria Limited.

[0068] Humanized antibody which is also called reshaped human antibody has been generated by transplanting the complementarity determining region (CDR) of antibody of a mammal other than the human, for example mouse antibody, into the CDR of human antibody.

The general recombinant DNA technology for preparation of such antibodies is also known (see European Patent Application EP 125023 and International Patent Publication WO 92-19759).

[0069] Specifically, a DNA sequence which was designed to ligate the CDR of mouse antibody with the framework region (FR) of human antibody is synthesized from several divided oligonucleotides having sections overlapping with one another at the ends thereof. The DNA thus obtained is ligated to the DNA encoding the C region of human antibody and then is integrated into an expression vector, which is introduced into a host for antibody production (see European Patent Application EP 239400 and International Patent Publication WO 92-19759).

[0070] For the FR of human antibody ligated through CDR, the complementarity determining region that forms a favorable antigen binding site is selected. When desired, amino acids in the framework region of the antibody variable region may be substituted so that the complementarity determining region of reshaped human antibody may form an appropriate antigen binding site (Sato, K. et al., Cancer Res. (1993) 53, 851-856).

[0071] For example, for chimeric antibody or humanized antibody, the C region of human antibody is used. As the C region of human antibody, there can be mentioned $\text{C}\gamma$, and, for example, $\text{C}\gamma 1$, $\text{C}\gamma 2$, $\text{C}\gamma 3$, and $\text{C}\gamma 4$ can be used. The C region of human antibody may be modified to improve the stability of antibody or the production thereof.

[0072] Chimeric antibody consists of the variable region of antibody derived from a mammal other than the human and the C region derived from human antibody, whereas humanized antibody consists of the complementarity determining region of antibody derived from a mammal other than the human and the framework region and the C region derived from human antibody. Accordingly, antigenicity thereof in the human body has been reduced so that they are useful as antibody for use in the present invention.

[0073] A preferred embodiment of the humanized antibody for use in the present invention includes humanized PM-1 antibody (see International Patent Publication WO 92-19759).

[0074] Antibody genes constructed as described above may be expressed and obtained in a known method. In the case of mammalian cells, expression may be accomplished using a vector containing a commonly used useful promoter, the antibody gene to be expressed, and DNA in which the poly A signal has been operably linked at 3' downstream thereof or a vector containing said DNA. Examples of the promoter/enhancer include human cytomegalovirus immediate early promoter/enhancer.

[0075] Additionally, as the promoter/enhancer which can be used for expression of antibody for use in the present invention, there are viral promoters/enhancers such as retrovirus, polyoma virus, adenovirus, and sim-

ian virus 40 (SV40), and promoters/enhancers derived from mammalian cells such as human elongation factor 1 α (HEF1 α).

[0076] For example, expression may be readily accomplished by the method of Mulligan et al. (Mulligan, R. C. et al., Nature (1979) 277, 108-114) when SV40 promoter/enhancer is used, or by the method of Mizushima et al. (Mizushima, S. and Nagata, S. Nucleic Acids Res. (1990) 18, 5322) when HEF1 α promoter/enhancer is used.

[0077] In the case of *E. coli*, expression may be conducted by operably linking a commonly used useful promoter, a signal sequence for antibody secretion, and the antibody gene to be expressed, followed by expression thereof. As the promoter, for example, there can be mentioned lacZ promoter and araB promoter. The method of Ward et al. (Ward, E.S. et al., Nature (1998) 341, 544-546; Ward, E.S. et al., FASEB J. (1992) 6, 2422-2427) may be used when lacZ promoter is used, and the method of Better et al. (Better, M. et al., Science (1988) 240, 1041-1043) may be used when araB promoter is used.

[0078] As the signal sequence for antibody secretion, when produced in the periplasm of *E. coli*, the pelB signal sequence (Lei, S. P. et al., J. Bacteriol. (1987) 169, 4379-4383) can be used. After separating the antibody produced in the periplasm, the structure of the antibody is appropriately refolded before use (see, for example, WO 96/30394).

[0079] As the origin of replication, there can be used those derived from SV40, polyoma virus, adenovirus, bovine papilloma virus (BPV) and the like. Furthermore, for the amplification of the gene copy number in the host cell system, expression vectors can include as selectable markers the aminoglycoside phosphotransferase (APH) gene, the thymidine kinase (TK) gene, *E. coli* xanthine guaninephosphoribosyl transferase (Ecogpt) gene, the dihydrofolate reductase (dhfr) gene and the like.

[0080] For the production of antibody for use in the present invention, any production system can be used. The production system of antibody preparation comprises the in vitro or the in vivo production system. As the in vitro production system, there can be mentioned a production system which employs eukaryotic cells and the production system which employs prokaryotic cells.

[0081] When the eukaryotic cells are used, there are the production systems which employ animal cells, plant cells, and fungal cells. Known animal cells include (1) mammalian cells such as CHO cells, COS cells, myeloma cells, baby hamster kidney (BHK) cells, HeLa cells, and Vero cells, (2) amphibian cells such as *Xenopus* oocytes, or (3) insect cells such as sf9, sf21, and Tn5. Known plant cells include, for example, those derived from *Nicotiana tabacum*, which may be subjected to callus culture. Known fungal cells include yeasts such as the genus *Saccharomyces*, more specifically *Saccharomyces cereviceae*, or filamentous fungi such as the ge-

nus Aspergillus, more specifically Aspergillus niger.

[0082] When the prokaryotic cells are used, there are the production systems which employ bacterial cells. Known bacterial cells include Escherichia coli (E. coli), and Bacillus subtilis.

[0083] By introducing via transformation the gene of the desired antibody into these cells and culturing the transformed cells in vitro, the antibody can be obtained. Culturing is conducted by a known method. For example, as the culture liquid, DMEM, MEM, RPMI1640, and IMDM can be used, and serum supplements such as fetal calf serum (FCS) may be used in combination. In addition, antibodies may be produced in vivo by implanting cells into which the antibody gene has been introduced into the abdominal cavity of an animal and the like.

[0084] As in vivo production systems, there can be mentioned those which employ animals and those which employ plants. When animals are used, there are the production systems which employ mammals and insects.

[0085] As mammals, goats, pigs, sheep, mice, and cattle can be used (Vicki Glaser, SPECTRUM Biotechnology Applications, 1993). Also as insects, silkworms can be used. When plants are used, tobacco, for example, can be used.

[0086] Antibody genes are introduced into these animals or plants, and the antibodies are produced in such animals or plants, and recovered. For example, an antibody gene is inserted into the middle of the gene encoding protein which is inherently produced in the milk such as goat β casein to prepare fusion genes. DNA fragments containing the fusion gene into which the antibody gene has been inserted are injected into a goat embryo, and the embryo is introduced into a female goat. The desired antibody is obtained from the milk produced by the transgenic goat borne to the goat who received the embryo or offspring thereof. In order to increase the amount of milk containing the desired antibody produced by the transgenic goat, hormones may be given to the transgenic goat as appropriate. (Ebert, K.M. et al., Bio/Technology (1994) 12, 699-702).

[0087] When silkworms are used, baculovirus into which the desired antibody gene has been inserted is infected to the silkworm, and the desired antibody can be obtained from the body fluid of the silkworm (Maeda, S. et al., Nature (1985) 315, 592-594). Moreover, when tobacco is used, the desired antibody gene is inserted into an expression vector for plants, for example pMON 530, and then the vector is introduced into a bacterium such as *Agrobacterium tumefaciens*. The bacterium is then infected to tobacco such as *Nicotiana tabacum* to obtain the desired antibody from the leaves of the tobacco (Julian, K.-C. Ma et al., Eur. J. Immunol. (1994) 24, 131-138).

[0088] When antibody is produced in in vitro or in vivo production systems, as described above, DNA encoding the heavy chain (H chain) or the light chain (L chain)

of antibody may be separately integrated into an expression vector and the hosts are transformed simultaneously, or DNA encoding the H chain and the L chain may be integrated into a single expression vector and the host is transformed therewith (see International Patent Publication WO 94-11523).

[0089] Antibodies for use in the present invention may be antibody fragments or modified versions thereof as long as they are preferably used. For example, as fragments of antibody, there may be mentioned Fab, $F(ab')_2$, Fv or single-chain Fv (scFv) in which Fv's of H chain and L chain were ligated via a suitable linker.

[0090] Specifically antibodies are treated with an enzyme, for example, papain or pepsin, to produce antibody fragments, or genes encoding these antibody fragments are constructed, and then introduced into an expression vector, which is expressed in a suitable host cell (see, for example, Co, M. S. et al., J. Immunol. (1994) 152, 2968-2976; Better, M. and Horwitz, A.H., Methods in Enzymology (1989) 178, 476-496; Plucktrun, A. and Skerra, A., Methods in Enzymology (1989) 178, 476-496; Lamoyi, E., Methods in Enzymology (1986) 121, 652-663; Rousseaux, J. et al., Methods in Enzymology (1986) 121, 663-669; Bird, R.E. et al., TIBTECH (1991) 9, 132-137).

[0091] scFv can be obtained by ligating the V region of H chain and the V region of L chain of antibody. In the scFv, the V region of H chain and the V region of L chain are preferably ligated via a linker, preferably a peptide linker (Huston, J.S. et al., Proc. Natl. Acad. Sci. USA (1988) 85, 5879-5883). The V region of H chain and the V region of L chain in the scFv may be derived from any of the above-mentioned antibodies. As the peptide linker for ligating the V regions, any single-chain peptide comprising, for example, 12-19 amino acid residues may be used.

[0092] DNA encoding scFv can be obtained using DNA encoding the H chain or the H chain V region of the above antibody and DNA encoding the L chain or the L chain V region of the above antibody as the template by amplifying the portion of the DNA encoding the desired amino acid sequence among the above sequences by the PCR technique with the primer pair specifying the both ends thereof, and by further amplifying the combination of DNA encoding the peptide linker portion and the primer pair which defines that both ends of said DNA are ligated to the H chain and the L chain, respectively.

[0093] Once DNAs encoding scFv are constructed, an expression vector containing them and a host transformed with said expression vector can be obtained by the conventional methods, and scFv can be obtained using the resultant host by the conventional methods.

[0094] These antibody fragments can be produced by obtaining the gene thereof in a similar manner to that mentioned above and by allowing it to be expressed in a host. "Antibody" as used in the claim of the present application encompasses these antibody fragments.

[0095] As modified antibodies, antibodies associated with various molecules such as polyethylene glycol (PEG) can be used. "Antibody" as used in the claim of the present application encompasses these modified antibodies. These modified antibodies can be obtained by chemically modifying the antibodies thus obtained. These methods have already been established in the art.

[0096] Antibodies produced and expressed as described above can be separated from the inside or outside of the host cell and then may be purified to homogeneity. Separation and purification of the antibody for use in the present invention may be accomplished by affinity chromatography. As the column used for such affinity chromatography, there can be mentioned Protein A column and Protein G column. Examples of the carriers used in the Protein A column are Hyper D, POROS, Sepharose F. F. and the like. Alternatively, methods for separation and purification conventionally used for proteins can be used without any limitation.

[0097] Separation and purification of the antibody for use in the present invention may be accomplished by combining, as appropriate, chromatography other than the above-mentioned affinity chromatography, filtration, ultrafiltration, salting-out, dialysis and the like. Chromatography includes, for example, ion exchange chromatography, hydrophobic chromatography, gel-filtration and the like. These chromatographies can be applied into high performance liquid chromatography (HPLC). Alternatively, reverse-phase HPLC can be used.

[0098] The concentration of antibody obtained in the above can be determined by the measurement of absorbance or by the enzyme-linked immunosorbent assay (ELISA) and the like. Thus, when absorbance measurement is employed, a sample is appropriately diluted with PBS(-) and then the absorbance is measured at 280 nm, followed by calculation using the absorption coefficient of 1.35 OD at 1 mg/ml. When the ELISA method is used, measurement is conducted as follows. Thus, 100 μ l of goat anti-human IgG (manufactured by TAGO) diluted to 1 μ g/ml in 0.1 M bicarbonate buffer, pH 9.6, is added to a 96-well plate (manufactured by Nunc), and is incubated overnight at 4 °C to immobilize the antibody. After blocking, 100 μ l each of appropriately diluted antibody of the present invention or a sample containing the antibody, or 100 μ l of human IgG (manufactured by CAPPEL) as the standard is added, and incubated at room temperature for 1 hour.

[0099] After washing, 100 μ l of 5000-fold diluted alkaline phosphatase-labeled anti-human IgG antibody (manufactured by BIO SOURCE) is added, and incubated at room temperature for 1 hour. After washing, the substrate solution is added and incubated, followed by the measurement of absorbance at 405 nm using the MICROPLATE READER Model 3550 (manufactured by Bio-Rad) to calculate the concentration of the desired antibody.

[0100] The altered IL-6 for use in the present invention

has an activity of binding to IL-6 receptor and does not transmit the biological activity of IL-6. Thus, the altered IL-6, though it competes with IL-6 receptor for binding to IL-6, does not transmit the biological activity of IL-6, and thereby it blocks signal transduction by IL-6.

[0101] Altered IL-6 may be constructed through the introduction of mutation by replacing amino acid residues of the amino acid sequence of IL-6. IL-6, the source of the altered IL-6, may be of any origin, but when the antigenicity is to be considered, it is preferably human IL-6.

[0102] Specifically, the secondary structure of IL-6 is predicted using a known molecular modeling program of the amino acid sequence, for example WHATIF (Vriend et al., J. Mol. Graphics (1990), 8, 52-56), and the overall effects on the amino acid residue to be replaced is evaluated. After an appropriate amino acid residue was determined, mutation is introduced by the commonly used polymerase chain reaction (PCR) method using a vector containing the base sequence encoding human IL-6 gene thereby to obtain a gene encoding altered IL-6. This is then integrated, as desired, into an appropriate expression vector, from which altered IL-6 can be obtained according to the expression, production and purification of said recombinant antibody.

[0103] Specific examples of the altered IL-6 are disclosed in Brakenhoff et al., J. Biol. Chem. (1994) 269, 86-93, and Savino et al., EMBO J. (1994) 13, 1357-1367, WO 96-18648, and WO 96-17869.

[0104] The IL-6 partial peptide or the IL-6 receptor partial peptide for use in the present invention has an activity of binding to IL-6 receptor or IL-6, respectively, and does not transmit the biological activity of IL-6. Thus, the IL-6 partial peptide or the IL-6 receptor partial peptide binds to IL-6 receptor or IL-6, respectively, and thereby capture it. As a result, they do not transmit the biological activity of IL-6, and block signal transduction of IL-6.

[0105] The IL-6 partial peptide or the IL-6 receptor partial peptide is a peptide comprising some or all of the amino acid sequence of the region involved in the binding to IL-6 and IL-6 receptor in the amino acid sequence of IL-6 or IL-6 receptor. Such a peptide generally comprises 10 - 80, preferably 20 - 50, and more preferably 20 - 40 amino acid residues.

[0106] The IL-6 partial peptide or the IL-6 receptor partial peptide can be constructed by specifying the region involved in the binding to IL-6 and IL-6 receptor in the amino acid sequence of IL-6 or IL-6 receptor, and by producing some or all of the amino acid sequence by a conventional method such as a genetic engineering technology or a peptide synthesis method.

[0107] In order to prepare the IL-6 partial peptide or the IL-6 receptor partial peptide by a genetic engineering technology, the DNA sequence encoding the desired peptide is integrated into an expression vector, from which the peptide can be obtained by the expression,

production, and purification of said recombinant antibody.

[0108] Preparation of the IL-6 partial peptide or the IL-6 receptor partial peptide by the peptide synthesis method can be effected using a method commonly used in peptide synthesis such as the solid phase synthesis or the liquid phase synthesis.

[0109] Specifically the method described in Zoku-Iyaku no Kaihatsu (Sequel to Development of Pharmaceuticals), Vol. 14, Peptide Gousei (Peptide Synthesis), edited by Haruaki Yajima, Hirokawa Shoten, 1991, may be used. The solid phase synthesis method used includes, for example, a reaction in which an amino acid corresponding to the C-terminal of the peptide to be synthesized is coupled to a support which is insoluble in organic solvents, and then an amino acid in which an α -amino group or a side chain functional group has been protected with an appropriate protecting group is condensed one amino acid at a time from the C-terminal to the N-terminal direction, and a reaction in which said protecting group of the α -amino group of the amino acid or the peptide coupled to the resin is eliminated is alternately repeated to elongate the peptide chain. The solid phase peptide synthesis methods are divided into the Boc method and the Fmoc method depending on the type of protecting group to be used.

[0110] After the synthesis of the desired peptide is complete, a deprotection reaction and a reaction for cleaving the peptide chain from the support is carried out. For cleavage from the peptide chain, hydrogen fluoride or trifluoromethanesulfonic acid in the Boc method and TFA in the Fmoc method are generally used. In the Boc method, for example, the above peptide resin is treated in hydrogen fluoride in the presence of anisole. Subsequently, the protecting group is eliminated and the peptide is recovered by cleaving from the support. By lyophilizing this, crude peptide can be obtained. On the other hand, in the Fmoc method, TFA, for example, is used in a manner similar to the above to effect the deprotection reaction and the cleavage reaction of the peptide from the support.

[0111] The crude peptide thus obtained can be applied to HPLC for its separation and purification. Its elution can be carried out in a water-acetonitrile solvent system that is commonly used for protein purification under an optimum condition. The fraction corresponding to the peak of the profile of the chromatography obtained is collected and lyophilized. The peptide fraction thus purified is identified by subjecting it to the analysis of molecular weight by mass spectroscopic analysis, the analysis of amino acid composition, or the analysis of amino acid sequence, and the like.

[0112] Specific examples of the IL-6 partial peptide or the IL-6 receptor partial peptide are disclosed in Japanese Unexamined Patent Publication (Kokai) 2(1990)-188600, Japanese Unexamined Patent Publication (Kokai) 7(1995)-324097, Japanese Unexamined Patent Publication (Kokai) 8(1996)-311098, and United States

Patent Publication US 5210075.

[0113] The activity of the IL-6 antagonist for use in the present invention can be evaluated using a conventionally known method. Specifically, the IL-6-dependent cell MH60.BSF2 is cultured, to which IL-6 is added, and the activity can be evaluated using the incorporation of ^3H -thymidine into the IL-6-dependent cell in the coexistence of the IL-6 antagonist. Alternatively, evaluation can be effected by culturing U266, an IL-6 receptor-expressing cell, adding thereto ^{125}I -labeled IL-6 and an IL-6 antagonist at the same time, and then by determining the ^{125}I -labeled IL-6 bound to the IL-6 receptor-expressing cell. In the above assay system, a negative control group containing no IL-6 antagonists, in addition to the group in which an IL-6 receptor antagonist is present, is set up, and the results obtained for them are compared to evaluate the IL-6-inhibiting activity of the IL-6 receptor antagonist.

[0114] In order to confirm the effects accomplished by the present invention, an IL-6 antagonist for use in the present invention is administered to animals that developed pancreatitis after an overdose of caerulein, and the effect of suppressing pancreatic edema and of improving the weight of the pancreas can be evaluated. As additional effects of the present invention, there are effects of preventing pancreatitis or the recurrence of pancreatitis.

[0115] The administration of caerulein to induce pancreatitis, for example, may be carried out according to the method described in the Example below. Animals in which pancreatitis is induced may be those commonly used in experiments such as mice and rats.

[0116] As described in the Example below, in the animals that developed pancreatitis, the administration of IL-6 receptor antibody resulted in suppression of the pancreas weight and improvement in the edema of the pancreas, and thus it was revealed that IL-6 antagonists such as anti-IL-6 receptor antibody exert a therapeutic effect on pancreatitis.

[0117] The subject to be treated in the present invention is mammals. The subject to be treated is preferably humans.

[0118] The preventive or therapeutic agents of the present invention may be administered, either orally or parenterally, systemically or locally. For example, intravenous injection such as drip infusion, intramuscular injection, intraperitoneal injection, subcutaneous injection, suppositories, intestinal lavage, oral enteric coated tablets, and the like can be selected, and the method of administration may be chosen, as appropriate, depending on the age and the conditions of the patient. The effective dosage is chosen from the range of 0.01 mg to 100 mg per kg of body weight per administration. Alternatively, the dosage in the range of 1 to 1000 mg, preferably 5 to 50 mg per patient may be chosen.

[0119] The preventive or therapeutic agents for pancreatitis of the present invention may contain pharmaceutically acceptable carriers or additives depending on

the route of administration. Examples of such carriers or additives include water, a pharmaceutical acceptable organic solvent, collagen, polyvinyl alcohol, polyvinylpyrrolidone, a carboxyvinyl polymer, carboxymethyl-cellulose sodium, polyacrylic sodium, sodium alginate, water-soluble dextran, carboxymethyl starch sodium, pectin, methyl cellulose, ethyl cellulose, xanthan gum, gum Arabic, casein, gelatin, agar, diglycerin, propylene glycol, polyethylene glycol, Vaseline, paraffin, stearyl alcohol, stearic acid, human serum albumin (HSA), mannitol, sorbitol, lactose, a pharmaceutically acceptable surfactant and the like. Additives used are chosen from, but are not limited to, the above or combinations thereof depending on the dosage form.

Examples

[0120] The present invention will now be explained in more detail with reference to the examples, reference examples, and experiments. It should be noted, however, that the present invention is not limited to them in any way.

Example 1.

[0121] To B6-hIL-6 transgenic mice or B6 mice (litter-mates of hIL-6 T transgenic mice) (Clinical Immunology and Immunopathology, 82: 117-124, 1997), caerulein (manufactured by Kyowa Hakko) dissolved in physiological saline at 50 µg/kg was intraperitoneally given every hour for 7 times. An anti-mouse IL-6 receptor monoclonal antibody M16-1 was administered to the mice at 1 mg/mouse via the tail vein immediately before caerulein administration.

[0122] As the control, the solvent (PBS) for the antibody was used. Eight hours later, the mice were euthanized to observe the weight of the pancreas, serum amylase, and the histology of the pancreas. The weight of the pancreas and body weight are shown in Figure 1. The histology of the pancreas is shown in Figures 2-4. Serum amylase was measured using the iodo-starch (blue starch) method. For the histology of the pancreas, a paraffin block of the pancreas was prepared and stained with hematoxylin eosin (HE) stain for microscopic observation.

[0123] In the IL-6 transgenic mice, the weight gain of the pancreas induced by caerulein administration was more pronounced as compared to the litter mate (normal mice).

[0124] Visual inspection also revealed that edema was more advanced as well as pancreatitis. By administering MR16-1 to the IL-6 transgenic mice, the weight gain of the pancreas induced by caerulein administration was suppressed. Histological comparison of the IL-6 transgenic mouse group (Figure 3) and the normal mouse group (Figure 2) has shown that areas of the edema etc. were smaller in the normal mice and the regions affected by neutrophil infiltration etc. were smaller.

This indicates that the administration of MR16-1 resulted in improvement (Figure 4). Thus, since the effect of anti-IL-6 receptor monoclonal antibody was observed in the mouse model of caerulein-induced acute pancreatitis, IL-6 antagonists such as anti-IL-6 receptor antibody are effective, by suppressing the effects of IL-6, in treatment, amelioration of severity, and prevention of onset of acute pancreatitis.

Example 2.

[0125] To B6-hIL-6 transgenic mice or B6 mice (litter-mates of hIL-6 T transgenic mice) (Clinical Immunology and Immunopathology, 82: 117-124, 1997), caerulein (manufactured by Kyowa Hakko) dissolved in physiological saline at 50 µg/kg was intraperitoneally given seven times at hourly intervals. In order to induce severe pancreatitis, LPS (lipopolysaccharide, manufactured by Sigma) was intraperitoneally given at 1 mg/ml concurrently with the initial caerulein administration. Anti-mouse IL-6 receptor monoclonal antibody M16-1 was administered to the mice at 1 mg/mouse via the tail vein 10 minutes before the initial caerulein administration. As the control, the solvent (PBS) for the antibody was used. Eight hours later, the mice were euthanized to measure the weight of the pancreas. The weight of the pancreas and body weight are shown in Figure 5.

[0126] In severe pancreatitis induced in the IL-6 transgenic mice, MR16-1 administration caused a pronounced improvement effect. Thus, in the mouse model of severe acute pancreatitis induced by caerulein and LPS, IL-6 antagonists such as anti-IL-6 receptor antibody are effective, by suppressing the effects of IL-6, in treatment, amelioration of severity, and prevention of onset of severe acute pancreatitis.

Reference example 1. Preparation of human soluble IL-6 receptor

[0127] Soluble IL-6 receptor was prepared by the PCR method using a plasmid pBSF2R.236 containing cDNA that encodes IL-6 receptor obtained according to the method of Yamasaki et al., (Yamasaki, K. et al., Science (1988) 241, 825-828). Plasmid pBSF2R.236 was digested with a restriction enzyme Sph I to obtain the cDNA of IL-6 receptor, which was then inserted into mpl8 (manufactured by Amersham). Using a synthetic oligoprimers designed to introduce a stop codon into the cDNA of IL-6 receptor, a mutation was introduced into the cDNA of IL-6 receptor by the PCR method using the in vitro Mutagenesis System (manufactured by Amersham). The procedure resulted in the introduction of a stop codon to the amino acid at position 345, and gave cDNA encoding soluble IL-6 receptor.

[0128] In order to express the cDNA of soluble IL-6 receptor in CHO cells, it was ligated to plasmid pSV (manufactured by Pharmacia) to obtain plasmid pSVL344. The cDNA of soluble IL-6 receptor that was

cleaved with Hind III-Sal I was inserted to plasmid pCEdhfr containing the cDNA of dhfr to obtain plasmid pCEdhfr344 that can be expressed in the CHO cells.

[0129] Ten µg of plasmid pCEdhfr344 was transfected to a dhfr-CHO cell line DXB-11 (Urlaub et al., Proc. Natl. Acad. Sci. USA (1980) 77, 4216-4220) by the calcium phosphate precipitation method (Chen C. et al., Mol. Cell. Biol. (1987) 7, 2745-2751). The transfected CHO cells were cultured for 3 weeks in a nucleoside-free α MEM selection medium containing 1 mM glutamine, 10% dialyzed FCS, 100 U/ml penicillin, and 100 µg/ml streptomycin.

[0130] The selected CHO cells were screened by the limiting dilution method to obtain a single CHO cell clone. The CHO cell clone was amplified in 20 nM - 200 nM methotrexate (MTX) to obtain a CHO cell line 5E27 that produces human soluble IL-6 receptor. The CHO cell line 5E27 was cultured in an Iscov-modified Dulbecco's medium (IMDM, manufactured by Gibco) containing 5% FBS. The culture supernatant was collected and the concentration of soluble IL-6 receptor in the culture supernatant was determined by ELISA. The result confirmed that soluble IL-6 receptor is present in the culture supernatant.

Reference example 2. Preparation of human IL-6 antibody

[0131] Ten µg of the recombinant IL-6 (Hirano et al., Immunol. Lett., 17:41, 1988) was immunized to BALB/c mice together with Freund's complete adjuvant, and this was repeated every week until anti-IL-6 antibody could be detected in the serum. Immune cells were extracted from local lymph node and were then fused with a myeloma cell line P3U1 using polyethylene glycol 1500. Hybridomas were selected according to the method of Oi et al. (Selective Methods in Cellular Immunology, W. H. Freeman and Co., San Francisco, 351, 1980) that employs the HAT medium, and the hybridoma that produces human IL-6 antibody was established.

[0132] The hybridoma that produces human IL-6 antibody was subjected to IL-6 binding assay as follows. Thus, a 96-well microtiter plate (manufactured by Dynatech Laboratories, Inc., Alexandria, VA) made of flexible polyvinyl was coated with 100 µl of goat anti-mouse Ig (10 µl/ml, manufactured by Cooper Biomedical, Inc., Malvern, PA) overnight at 4 °C. Subsequently, the plate was treated with 100 µl of PBS containing 1% bovine serum albumin (BSA) at room temperature for 2 hours.

[0133] After washing it in PBS, 100 µl of the hybridoma culture supernatant was added to each well, and then was incubated overnight at 4 °C. The plate was washed, ¹²⁵I-labeled recombinant IL-6 was added to each well to a concentration of 2000 cpm/0.5 ng/well, and then radioactivity of each well after washing was determined by a gamma counter (Beckman Gamma 9000, Beckman Instruments, Fullerton, CA). Of 216 hybridoma clones, 32 were positive in the IL-6 binding as-

say. From these clones, stable MH166.BSF2 was finally obtained. Anti-IL-6 antibody MH166 produced by said hybridoma has a subtype of IgG1 κ.

[0134] Then, the IL-6-dependent mouse hybridoma clone MH60.BSF2 was used to examine a neutralizing activity with respect to the growth of the hybridoma by MH166 antibody. MH60.BSF2 cells were dispensed to 1×10^4 /200 µl/well, and samples containing MH166 antibody were added thereto, cultured for 48 hours, 0.5 µCi/well of ³H-thymidine (New England Nuclear, Boston, MA) was added, and the culturing was continued for further 6 hours. The cells were placed on a glass filter paper and were treated by the automatic harvester (Labo Mash Science Co., Tokyo, Japan). As the control, rabbit anti-IL-6 antibody was used.

[0135] As a result, MH166 antibody inhibited, in a dose dependent manner, the incorporation of ³H-thymidine of MH60.BSF2 cells induced by IL-6. This revealed that MH166 antibody neutralizes the activity of IL-6.

Reference example 3. Preparation of human anti-IL-6 receptor antibody

[0136] Anti-IL-6 receptor antibody MT18 prepared by the method of Hirata et al. (Hirata, Y. et al. J. Immunol., 143, 2900-2906, 1989) was bound to CNBr-activated Sepharose 4B (manufactured by Pharmacia Fine Chemicals, Piscataway, NJ) according to the attached regimen, and IL-6 receptor (Yamasaki, K. et al., Science (1988) 241, 825-828) was purified. A human myeloma cell line U266 was solubilized with 1 mM p-para-aminophenyl methane sulfonyl fluoride hydrochloride (manufactured by Wako Chemicals) containing 1% digitonin (manufactured by Wako Chemicals), 10 mM triethanolamine (pH 7.8) and 0.15 M NaCl (digitonin buffer), and mixed with MT18 antibody bound to Sepharose 4B beads. Then, the beads were washed six times with the digitonin buffer to prepare the partially purified IL-6 receptor for use in immunization.

[0137] BALB/c mice were immunized four times, at ten-day intervals, with the above partially purified IL-6 receptor obtained from 3×10^9 U266 cells, and then a hybridoma was prepared using a standard method. The hybridoma culture supernatant from the growth-positive well was tested for its activity of binding to IL-6 receptor according to the method described below. 5×10^7 U266 cells were labeled with ³⁵S-methionine (2.5 mCi) and were solubilized with the above digitonin buffer. The solubilized U266 cells were mixed with a 0.04 ml volume of MT18 antibody bound to Sepharose 4B beads, and then were washed six times with the digitonin buffer. ³⁵S-methionine-labeled IL-6 receptor was eluted with 0.25 ml of the digitonin buffer (pH 3.4) and was neutralized in 0.025 ml of 1M Tris (pH 7.4).

[0138] 0.05 ml of the hybridoma culture supernatant was mixed with 0.01 ml of Protein G Sepharose (manufactured by Pharmacia). After washing, Sepharose was incubated with 0.005 ml of ³⁵S-labeled IL-6 receptor so-

lution prepared as described above. The immunoprecipitate was analyzed by SDS-PAGE to determine the hybridoma culture supernatant that reacts with IL-6 receptor. As a result, the reaction-positive hybridoma clone PM-1 was established. The antibody produced from the hybridoma PM-1 has a subtype of IgG1κ.

[0139] The inhibitory activity of IL-6 binding of the antibody produced by the hybridoma PM-1 to human IL-6 receptor was studied using the human myeloma cell line U266. A human recombinant IL-6 was prepared from *E. coli* (Hirano et al., Immunol. Lett., 17:41-45, 1988), and was labeled with ¹²⁵I using the Bolton-Hunter reagent (New England Nuclear, Boston, MA) (Taga, T. et al., J. Exp. Med. (1987) 166, 967-981). 4 x 10⁵ U266 cells were cultured with the culture supernatant of 70% (v/v) hybridoma PM-1 together with 14,000 cpm of ¹²⁵I-labeled IL-6 for one hour. Seventy μl of the sample was layered on 300 μl FCS in a 400 μl microfuge polyethylene tube. After centrifugation, the radioactivity of the cell was determined.

[0140] The result revealed that the antibody produced by the hybridoma PM-1 inhibits the binding of IL-6 to IL-6 receptor.

Reference example 4. Preparation of mouse anti-IL-6 receptor antibody

[0141] A monoclonal antibody directed against mouse IL-6 receptor was prepared according to the method described in Saito, et al., J. Immunol. (1993) 147, 168-173.

[0142] The CHO cells that produce mouse soluble IL-6 receptor were cultured in the IMDM culture liquid containing 10% FCS. From the culture supernatant, mouse soluble IL-6 receptor was purified using mouse soluble IL-6 receptor antibody RS12 (see Saito, et al., supra) and an affinity column fixed to Affigel 10 gel (manufactured by Biorad).

[0143] The mouse soluble IL-6 receptor (50 μg) thus obtained was mixed with Freund's complete adjuvant, which was then injected to the abdomen of Wistar rats. From 2 weeks after the administration, the animals were boosted with Freund's incomplete adjuvant. On day 45, the rat spleen cells were collected, and the cells at about 2 x 10⁸ were fused with 1 x 10⁷ mouse myeloma cells P3U1 using a 50% PEG1500 (manufactured by Boehringer Mannheim) according to the conventional method, and then were screened by the HAT culture medium.

[0144] After the culture supernatant was added to the plate coated with rabbit anti-rat IgG antibody (manufactured by Cappel), mouse soluble IL-6 receptor was reacted. Subsequently, using rabbit anti-mouse IL-6 receptor antibody and alkaline phosphatase-labeled sheep anti-rabbit IgG, hybridomas producing antibody directed against mouse soluble IL-6 receptor were screened by ELISA. After antibody production was confirmed, the hybridoma clones were subscreened twice to obtain a single hybridoma clone. The clone was des-

ignated as MR16-1.

[0145] The neutralizing activity of the antibody produced by the hybridoma on signal transduction of mouse IL-6 was examined by ³H-thymidine incorporation using MH60.BSF2 cells (Matsuda, T. et al., J. Immunol. (1988) 18, 951-956). To a 96-well plate, MH60.BSF2 cells were prepared at 1 x 10⁴ cells/200 μl/well. To the plate were added 10 pg/ml mouse IL-6 and MR16-1 antibody or RS12 antibody at 12.3 - 1000 ng/ml, and then were cultured at 37°C and 5% CO₂ for 44 hours and then 1 μCi/well of ³H-thymidine was added. After 4 hours, the incorporation of ³H-thymidine was measured. As a result, MR16-1 antibody suppressed the incorporation of ³H-thymidine of the MH60.BSF2 cells.

[0146] Thus, it was demonstrated that the antibody produced by the hybridoma MR16-1 inhibits the binding of IL-6 to IL-6 receptor.

Industrial Applicability

[0147] In accordance with the present invention, it was shown that IL-6 antagonists such as anti-IL-6 receptor antibody have a therapeutic effect on pancreatitis. Thus, it was demonstrated that IL-6 antagonists are useful as a therapeutic agent for acute pancreatitis and the like.

Claims

1. A preventive or therapeutic agent for pancreatitis comprising an interleukin-6 (IL-6) antagonist as an active ingredient.
2. The preventive or therapeutic agent according to claim 1 in which the IL-6 antagonist is an antibody directed against IL-6 receptor.
3. The preventive or therapeutic agent according to claim 2 in which the antibody directed against IL-6 receptor is a monoclonal antibody directed against IL-6 receptor.
4. The preventive or therapeutic agent according to claim 3 in which the antibody directed against IL-6 receptor is a monoclonal antibody directed against human IL-6 receptor.
5. The preventive or therapeutic agent according to claim 3 in which the antibody directed against IL-6 receptor is a monoclonal antibody directed against mouse IL-6 receptor.
6. The preventive or therapeutic agent according to any of claims 2 to 5 in which the antibody directed against IL-6 receptor is a recombinant antibody.

7. The preventive or therapeutic agent according to claim 4 in which the monoclonal antibody directed against human IL-6 receptor is PM-1 antibody.
8. The preventive or therapeutic agent according to claim 5 in which the monoclonal antibody directed against mouse IL-6 receptor is MR16-1 antibody.
9. The preventive or therapeutic agent according to any of claims 2 to 4 in which the antibody directed against IL-6 receptor is a chimeric antibody or a humanized antibody directed against IL-6 receptor.
10. The preventive or therapeutic agent according to claim 9 in which the humanized antibody directed against IL-6 receptor is a humanized PM-1 antibody.
11. The preventive or therapeutic agent according to any of claims 1 to 10 in which the pancreatitis is acute pancreatitis.
12. An agent for suppressing pancreatic edema said agent comprising an IL-6 antagonist as an active ingredient.
13. An agent for suppressing pancreatic edema said agent comprising an antibody directed against IL-6 receptor as an active ingredient.
14. A method of preventing or treating pancreatitis which method comprises administering an interleukin-6 (IL-6) antagonist at an amount effective to prevent or treat pancreatitis to the subject in need of such prevention or treatment.
15. The preventive or therapeutic method according to claim 14 in which the IL-6 antagonist is an antibody directed against IL-6 receptor.
16. The preventive or therapeutic method according to claim 15 in which the antibody directed against IL-6 receptor is a monoclonal antibody directed against IL-6 receptor.
17. The preventive or therapeutic method according to claim 16 in which the antibody directed against IL-6 receptor is a monoclonal antibody directed against human IL-6 receptor.
18. The preventive or therapeutic method according to claim 16 in which the antibody directed against IL-6 receptor is a monoclonal antibody directed against mouse IL-6 receptor.
19. The preventive or therapeutic method according to claim 15 in which the antibody directed against IL-6 receptor is a recombinant antibody.
20. The preventive or therapeutic method according to claim 17 in which the monoclonal antibody directed against human IL-6 receptor is PM-1 antibody.
21. The preventive or therapeutic method according to claim 18 in which the monoclonal antibody directed against mouse IL-6 receptor is MR16-1 antibody.
22. The preventive or therapeutic method according to claim 15 in which the antibody directed against IL-6 receptor is a chimeric antibody or a humanized antibody directed against IL-6 receptor.
23. The preventive or therapeutic method according to claim 22 in which the humanized antibody directed against IL-6 receptor is a humanized PM-1 antibody.
24. The preventive or therapeutic method according to claim 14 in which the pancreatitis is acute pancreatitis.
25. A method of suppressing pancreatic edema in pancreatitis which method comprises administering an IL-6 antagonist to the subject.
26. A method of suppressing pancreatic edema which method comprises administering an antibody directed against IL-6 receptor at an amount effective to prevent or treat pancreatic edema to the subject in need of suppressing pancreatic edema.
27. The use of an interleukin-6 (IL-6) antagonist for the production of a preventive or therapeutic agent for pancreatitis.
28. The use according to claim 27 in which the IL-6 antagonist is an antibody directed against IL-6 receptor.
29. The use according to claim 28 in which the antibody directed against IL-6 receptor is a monoclonal antibody directed against IL-6 receptor.
30. The use according to claim 29 in which the antibody directed against IL-6 receptor is a monoclonal antibody directed against human IL-6 receptor.
31. The use according to claim 29 in which the antibody directed against IL-6 receptor is a monoclonal antibody directed against mouse IL-6 receptor.
32. The use according to any of claims 28 to 31 in which the antibody directed against IL-6 receptor is a recombinant antibody.
33. The use according to claim 30 in which the monoclonal antibody directed against human IL-6 receptor is PM-1 antibody.

tor is PM-1 antibody.

34. The use according to claim 31 in which the monoclonal antibody directed against mouse IL-6 receptor is MR16-1 antibody. 5
35. The use according to any of claims 28 to 30 in which the antibody directed against IL-6 receptor is a chimeric antibody or a humanized antibody directed against IL-6 receptor. 10
36. The use according to claim 35 in which the humanized antibody directed against IL-6 receptor is a humanized PM-1 antibody. 15
37. The use according to any of claims 27 to 36 in which the pancreatitis is acute pancreatitis.
38. The use of an IL-6 antagonist for the production of an agent for suppressing pancreatic edema in pancreatitis. 20
39. The use of antibody directed against IL-6 receptor for the production of an agent for suppressing pancreatic edema in pancreatitis. 25

30

35

40

45

50

55

Fig.1




Fig. 2




Fig. 3




Fig. 4





Fig. 5



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/04533

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl. ⁶ A61K45/00, A61K39/395		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl. ⁶ A61K45/00, A61K39/395		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CA (STN), REGISTRY (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X Y	Chemical Abstracts, Vol. 121, Abstract 154705 (1994), "Islet inflammation and hyperplasia induced by the pancreatic islet-specific overexpression of interleukin-6 in transgenic mice"	1, 27 2-13, 28-39
X Y	H.Hank Simms et al, "Polymorphonuclear Leukocyte Dysregulation During the Systemic Inflammatory Response Syndrome", Blood, Vol. 83, No.5 (1994) P.1398-1407	1, 27 2-13, 28-39
Y	EP, 572118, A1 (TOSOH CORP.) 01 December, 1993 (01.12.93), & JP, 5-304986, A & US, 5866689, A & DE, 69319662, E & IL, 105541, A	1-4, 6, 10-13 27-30, 32, 36-39
Y	WO, 92/19759, A1 (CHUGAI SEIYAKU KABUSHIKI KAISHA), 12 November, 1992 (12.11.92), & EP, 628639, A1 & DE, 69229482, E & US, 5795965, A & JP, 5-227970, A	1-13, 27-39
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&" document member of the same patent family</p>		
Date of the actual completion of the international search 16 November, 1999 (16.11.99)		Date of mailing of the international search report 30 November, 1999 (30.11.99)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/04533

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	Fumimaro Takaku et al. "The Merck Manual 16 th Ed." (Publication on 01 May, 1995 3 rd printing) P.760-766, Column " Pancreatitis"	12,13,38,39

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/04533

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 14 - 26
because they relate to subject matter not required to be searched by this Authority, namely:

The subject matter of claims 14 - 26 relates to a method for treatment of the human body by therapy, which does not require an international search report by this International Search Authority in accordance with PCT Article 17(2) (a)(i) and Rule 39.1(iv).
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1992)